

Table 1 AAC, TPC, TFC and TAC of 15 kiwifruit cultivars (Adopted from Zhang et al., 2020)

Cultivar	AAC (mg/100 g)	TPC (mg GAE/100 g)	TFC (mg CTE/100 g)	TAC (mg CGE/100 g)
Cuixiang	145.25 ± 0.75 h	131.03 ± 2.84 e	20.39 ± 3.55 de	-
Hayward	92.75 ± 1.75 l	78.04 ± 3.27 h	10.25 ± 2.13 f	-
Xuxiang	105.42 ± 0.63 k	85.49 ± 2.55 g	12.56 ± 3.51 ef	-
Guichang	165.27 ± 3.25 e	133.94 ± 3.52 e	5.69 ± 2.68 f	-
Ruiyu	124.49 ± 0.89 j	110.57 ± 3.46 f	12.00 ± 1.47 ef	-
Jinhong No.1	52.39 ± 1.88 n	80.09 ± 1.91 h	22.19 ± 3.29 d	0.52 ± 0.07 c
Hongshi No.2	248.16 ± 2.67 a	216.37 ± 0.73 a	50.89 ± 3.02 a	3.65 ± 0.27 a
Hongyang	154.57 ± 1.72 g	129.75 ± 1.59 e	40.69 ± 3.01 b	1.02 ± 0.02 b
Jinhong 50	161.04 ± 1.70 f	142.89 ± 3.78 c	31.64 ± 3.69 c	0.72 ± 0.00 c
Oriental Red	181.07 ± 2.10 d	144.53 ± 0.77 c	23.72 ± 5.71 d	1.58 ± 0.13 b
Huayou	187.33 ± 1.88 c	146.46 ± 2.43 c	23.58 ± 1.44 d	-
Sungold	137.42 ± 1.38 i	113.09 ± 3.22 f	16.33 ± 3.06 e	-
Jinyan	73.83 ± 1.28 m	74.06 ± 0.97 h	15.17 ± 1.12 ef	-
Jintao	151.74 ± 1.61 g	141.7 ± 2.19 d	14.69 ± 4.52 ef	-
Puyu	233.99 ± 1.61 b	158.66 ± 2.35 b	19.61 ± 3.26 d	-

Note: The data are given as the mean ± SD ($n = 3$). Mean values in each column with unlike letters are significantly different among cultivars ($p < 0.05$) (Adopted from Zhang et al., 2020)