

Table 1 Effect of foliar spray of different concentrations of boron on plant height of cauliflower (*Brassica oleracea* var. *botrytis*) at Marin, Sindhuli, Nepal (2024)

Treatment	Plant height (cm)			
	45 DAT	60 DAT	75 DAT	80 DAT
Control	32.16	46.47	52.45	53.32
0.025% Boron (1-time spray)	35.21	49.26	58.15	58.44
0.025% Boron (2- times spray)	29.62	48.65	53.52	54.44
0.025% Boron (3- times spray)	33.99	49.69	56.09	56.24
0.05% Boron (1-time spray)	34.96	49.13	57.70	59.88
0.05% Boron (2- times spray)	31.22	46.88	55.73	59.26
0.05% Boron (3- times spray)	34.01	48.72	56.49	58.05
0.075% Boron (1-time spray)	34.30	47.73	52.56	54.05
0.075% Boron (2- times spray)	34.02	51.63	56.93	59.63
0.075% Boron (3- times spray)	33.08	49.01	57.40	57.63
SEm ($\pm$)	1.338	1.87	2.95	2.53
LSD (0.05)	3.97	5.57	8.77	7.54
F test	NS	NS	NS	NS
CV (%)	6.97	6.66	9.18	7.73
Grand mean	33.26	48.72	55.70	56.85

Note: LSD=Least significant difference, SEM=Standard error of mean, CV=Coefficient of variation, DAT=Days after transplanting, NS=Non-significant