

Table 2 Effect of treatments on primary branches

Treatment	Primary branches of different varieties of okra on different days				
	30 DAS	45 DAS	60 DAS	75 DAS	90 DAS
Arka Anamika	1.40 ^b	1.80 ^c	2.30 ^c	2.40 ^b	2.00 ^a
Rukmani	1.30 ^b	1.65 ^{bc}	2.10 ^{bc}	2.30 ^b	2.55 ^d
Raja	1.05 ^a	1.30 ^a	1.50 ^a	1.75 ^a	2.15 ^b
Special-45	1.30 ^b	1.50 ^{ab}	1.95 ^b	2.20 ^b	2.35 ^a
Arzine	1.45 ^b	1.70 ^{bc}	2.10 ^{bc}	2.75 ^c	3.00 ^e
LSD(0.05)	0.21	0.22	0.31	0.14	0.11
F-test	*	**	**	***	***
SEm(±)	0.06	0.07	0.10	0.09	0.53
CV(%)	10.6	9.2	10.2	8.5	3.1
Grand Mean	1.30	1.59	2.00	2.21	2.41

Note: *Significant at 5% level of significance, **Significant at 1% level of significance, ***Significant at 0.1% level of significance, LSD: least significant difference, SEm: Standard error of the mean, CV: coefficient of difference, DAS: Days After Sowing