

Table 3 Effect of foliar spray of different concentrations of boron on leaf length of cauliflower (*Brassica oleracea* var. *botrytis*) at Marin, Sindhuli, Nepal (2024)

Treatment	Leaf length (cm)			
	45 DAT	60 DAT	75 DAT	80 DAT
Control	21.27	32.05	38.98	41.47
0.025% Boron (1-time spray)	23.51	34.30	45.34	46.55
0.025% Boron (2- times spray)	20.17	33.87	38.51	39.17
0.025% Boron (3- times spray)	23.11	35.18	41.37	42.66
0.05% Boron (1-time spray)	23.06	33.86	42.66	47.35
0.05% Boron (2- times spray)	19.62	32.24	40.48	43.95
0.05% Boron (3- times spray)	21.04	32.20	43.03	38.37
0.075% Boron (1-time spray)	22.73	34.14	39.85	43.61
0.075% Boron (2- times spray)	23.57	34.28	40.61	42.48
0.075% Boron (3- times spray)	20.41	33.75	40.25	46.02
SEm ($\pm$)	1.01	1.22	2.00	3.68
LSD (0.05)	2.99	3.62	5.94	10.94
F test	NS	NS	NS	NS
CV (%)	7.98	6.29	8.43	14.77
Grand Mean	21.85	33.59	41.11	43.16

Note: LSD=Least Significant Difference, SEM=Standard Error of Mean, CV=Coefficient of Variation, DAT=Days After Transplanting, NS=Non-significant