Table 4 Interaction effect of varieties and mlching method on different parameters of potato

Treatment	Germination (%)	Height (cm)	Tuber weight (kg/m ²)	Tubers no (>100gm)
Khumal Seto*Black plastic mulch	93.33 ^a	40.87^{a}	5.10 ^a	12.67 ^a
Khumal Seto*silver plastic mulch	90.67 ^a	36.23 ^b	4.3 ^b	8.33 ^b
Khumal Seto*Organic mulch	82.67 ^a	34.3 ^{bc}	3.35°	8.67 ^b
Khumal Seto*Control	86.67 ^a	31.87^{cd}	2.19 ^d	3.33°
Cardinal*Black plastic mulch	90.67 ^a	28.77^{d}	2.16 ^d	1 ^{cd}
Cardinal*silver plastic mulch	76^{a}	31.16^{cd}	1.88 ^d	0.33^{d}
Cardinal*Organic mulch	48.67 ^b	22.56e	1.59 ^{de}	1.33 ^{cd}
Cardinal*Control	30^{c}	17.93 ^f	0.98e	0^{d}
LSD (0.05)	2.67	3.86	0.67	2.67
SEM (±)	5.61	1.27	0.22	0.88
F probability	< 0.01	< 0.05	< 0.01	< 0.01

Note: \pm Means of 3 replications. Means in column with same superscript is not significantly different by DMRT at 0.05 level, LSD: Least Significant Difference, CV: Coefficient of variation, SEM (\pm): Standard error of mean