

Table 4 Canopy Spread (cm) of eight strawberry varieties evaluated under low-chilling conditions in Chitwan, Nepal (2020-2021)

Treatments	Canopy spread (cm)						
	30 DAT	45 DAT	60 DAT	75 DAT	90 DAT	105 DAT	120 DAT
Florida Festival	19.06 ^{ab}	27.39 ^a	37.62 ^a	44.63 ^a	48.54 ^a	50.13 ^a	51.36 ^a
Florida Beauty	17.05 ^{bc}	20.20 ^c	27.13 ^c	30.95 ^c	33.33 ^d	34.52 ^d	35.29 ^d
Sweet Sensation	17.51 ^{bc}	27.84 ^a	37.96 ^a	45.54 ^a	48.29 ^a	49.67 ^{ab}	50.62 ^a
Winter Star	17.24 ^{bc}	25.26 ^{ab}	32.38 ^b	40.34 ^b	43.16 ^b	45.02 ^{bc}	46.02 ^b
Elyana	17.69 ^{abc}	23.94 ^b	31.35 ^b	38.04 ^b	40.41 ^c	41.50 ^c	42.22 ^c
Ruby Gem	19.90 ^a	27.41 ^a	36.26 ^a	44.60 ^a	47.91 ^a	49.22 ^{ab}	50.17 ^a
Winter Dawn	15.91 ^c	19.28 ^c	25.70 ^{cd}	31.06 ^c	33.50 ^d	34.81 ^d	35.51 ^d
Nyho	13.51 ^d	17.50 ^c	23.24 ^d	28.94 ^c	31.48 ^d	28.45 ^e	32.92 ^d
Grand Mean	17.24	23.60	31.45	38.01	40.83	41.66	43.01
SEM	0.69	1.45	2.01	2.43	2.56	2.92	2.70
LSD	2.15***	2.66***	2.84***	2.90***	2.61***	4.66***	2.54***
CV	8.47	7.62	6.15	5.21	4.36	7.61	4.02

Note: In the DMRT, the significance level is as follows: ns=Not significant at 5% ($p \leq 0.05$). *=Significant at 5% ($p \leq 0.01$),

=Significant at 1% ($p \leq 0.001$), *=Significant at less than 1% ($p < 0.001$). SEM: Standard error of the mean, LSD: Least significant difference, CV: Coefficient of variation