

Table 4 Effect of foliar spray of different concentrations of boron on leaf breadth of cauliflower (*Brassica oleracea* var. *botrytis*) at Marin, Sindhuli, Nepal (2024)

Treatment	Leaf breadth (cm)			
	45 DAT	60 DAT	75 DAT	80 DAT
Control	13.42	20.60	18.08	19.44
0.025% Boron (1-time spray)	15.07	22.63	21.05	20.06
0.025% Boron (2- times spray)	13.05	20.81	18.99	18.30
0.025% Boron (3- times spray)	14.70	33.57	20.07	21.10
0.05% Boron (1-time spray)	15.35	22.50	20.83	22.33
0.05% Boron (2- times spray)	13.48	20.56	19.66	21.01
0.05% Boron (3- times spray)	14.14	21.60	20.84	21.14
0.075% Boron (1-time spray)	15.59	21.46	18.98	19.45
0.075% Boron (2- times spray)	16.78	22.91	20.98	22.05
0.075% Boron (3- times spray)	14.76	22.54	21.02	21.94
SEm ($\pm$)	0.79	0.75	1.12	1.324
LSD (0.05)	2.35	2.25	3.35	3.93
F test	NS	NS	NS	NS
CV (%)	9.39	8.97	9.73	11.08
Grand mean	14.63	14.63	20.05	20.68

Note: LSD=Least Significant Difference, SEM=Standard Error of Mean, CV=Coefficient of Variation, DAT=Days After Transplanting, NS=Non-significant