

Table 6 Effect of foliar spray of different concentrations of boron on length diameter of the hollowness of cauliflower (*Brassica oleracea* var. *botrytis*) at Marin, Sindhuli, Nepal (2024)

Treatment	Length of hollowness	Hollowness of diameter
Control	8.57 ^a	1.83 ^a
0.025% Boron (1-time spray)	4.57 ^b	1.30 ^{ab}
0.025% Boron (2- times spray)	3.83 ^{bc}	1.16 ^{abc}
0.025% Boron (3- times spray)	0.41 ^c	0.16 ^d
0.05% Boron (1-time spray)	4.84 ^b	1.40 ^{ab}
0.05% Boron (2- times spray)	2.24 ^{bc}	0.76 ^{bcd}
0.05% Boron (3- times spray)	0.22 ^c	0.43 ^{bcd}
0.075% Boron (1-time spray)	0.66 ^c	0.60 ^{bcd}
0.075% Boron (2- times spray)	1.15 ^{bc}	0.43 ^{bcd}
0.075% Boron (3- times spray)	0.16 ^c	0.20 ^{cd}
SEm (\pm)	1.18	0.29
LSD (0.05)	3.50	0.88
F test	**	**
CV (%)	76.56	61.89
Grand Mean	2.67	0.83

Note: Means followed by common letter(s) within column are non-significantly different based on DMRT at P=0.05; LSD=Least Significant Difference, SEM=Standard Error of Mean, CV=Coefficient of Variation, **=significant at 1% level of significance